

# Inapproximability of Maximum $r$ -Regular Induced Connected Subgraph Problems

Yuichi Asahiro<sup>1</sup>, Hiroshi Eto<sup>2</sup>, and Eiji Miyano<sup>2</sup>

<sup>1</sup>Department of Information Science, Kyushu Sangyo University,  
Fukuoka 813-8503, Japan. asahiro@is.kyusan-u.ac.jp

<sup>2</sup>Department of System Design and Informatics, Kyushu Institute of Technology,  
Fukuoka 820-8502, Japan. {eto@theory., miyano@}ces.kyutech.ac.jp

**Abstract**—Given a graph  $G = (V, E)$  on  $n$  vertices, the MAXIMUM  $r$ -REGULAR INDUCED CONNECTED SUBGRAPH ( $r$ -MaxRICS) problems ask for a maximum sized subset of vertices  $S \subseteq V$  such that the induced subgraph  $G[S]$  on  $S$  is connected and  $r$ -regular. For  $r = 2$ , it is known that 2-MaxRICS is  $\mathcal{NP}$ -hard and cannot be approximated within a factor of  $n^{1-\varepsilon}$  in polynomial time for any  $\varepsilon > 0$  if  $\mathcal{P} \neq \mathcal{NP}$ . In this paper, we show that  $r$ -MaxRICS is  $\mathcal{NP}$ -hard for any fixed integer  $r \geq 3$ , and furthermore  $r$ -MaxRICS cannot be approximated within a factor of  $n^{1/6-\varepsilon}$  in polynomial time for any  $\varepsilon > 0$  if  $\mathcal{P} \neq \mathcal{NP}$ .

**Keywords:** induced connected subgraph, regularity,  $\mathcal{NP}$ -hardness, inapproximability

## 1. Introduction

In this paper we only consider simple undirected graphs with no loops and no multiple edges. Let  $G = (V(G), E(G))$  be a graph, where  $V(G)$  and  $E(G)$  denote the set of vertices and the set of edges in  $G$ , respectively. A graph  $G_S$  is a subgraph of a graph  $G$  if  $V(G_S) \subseteq V(G)$  and  $E(G_S) \subseteq E(G)$ . For a subset of vertices  $S \subseteq V(G)$ , by  $G[S]$ , we mean the subgraph of  $G$  induced on  $S$ , which is called the *induced subgraph*.

The problem MAXIMUM INDUCED SUBGRAPH (MaxIS) of finding the maximum number of vertices that induces a subgraph satisfying some properties is one of the most fundamental problems in the fields of graph theory and combinatorial optimization, and thus extensively studied in these decades. Unfortunately, however, it is well known that the MaxIS problem is  $\mathcal{NP}$ -hard for a large class of interesting properties. For example, in [7], Lund and Yannakakis prove that the MAXIMUM INDUCED SUBGRAPH problem for the natural properties such as *acyclicity*, *planarity*, and *bipartiteness* cannot be approximated within a factor of  $n^{1-\varepsilon}$  in polynomial time for any positive constant  $\varepsilon$  if  $\mathcal{P} \neq \mathcal{NP}$ , where  $n$  is the number of the vertices in the input graph.

## 1.1 Our Problems and Results

A graph is  $r$ -regular if the degree of every vertex is exactly  $r$ . The *regularity* of graphs must be one of the most basic properties. In this paper we consider the following variant of the MaxIS problem, i.e., the desired properties the induced subgraph must satisfy are *regularity* and *connectivity*:

MAXIMUM  $r$ -REGULAR INDUCED CONNECTED SUBGRAPH ( $r$ -MaxRICS)

Input: A graph  $G = (V, E)$  and an integer  $r$ .

Goal: Find a maximum subset of vertices  $S \subseteq V$  such that the induced subgraph  $G[S]$  on  $S$  is connected and  $r$ -regular.

Since a *clique* is connected and regular, the MAXIMUM CLIQUE problem may be regarded as a special one of  $r$ -MaxRICS. The MAXIMUM CLIQUE is very difficult even to approximate [5]. Clearly, however, the problem of finding a clique of a constant degree is solvable in polynomial time. On the other hand,  $r$ -MaxRICS is hard even if  $r$  is a small constant as follows: The problem 2-MaxRICS is known as LONGEST INDUCED CYCLE problem since a 2-regular subgraph means a cycle in the input graph. In [6] Kann shows the following inapproximability for 2-MaxRICS:

**Theorem 1 ([6]):** 2-MaxRICS cannot be approximated in polynomial time within a factor of  $n^{1-\varepsilon}$  for any constant  $\varepsilon > 0$  if  $\mathcal{P} \neq \mathcal{NP}$ , where  $n$  is the number of vertices in the input graph.

In [3] Bonifaci, Di Iorio, and Laura consider the following problem and show its  $\mathcal{NP}$ -hardness:

MAXIMUM REGULAR INDUCED SUBGRAPH (MaxRIS)

Input: A graph  $G = (V, E)$ .

Goal: Find a maximum subset of vertices  $S \subseteq V$  such that the induced subgraph  $G[S]$  on  $S$  is regular.

Strictly speaking, MaxRIS is slightly different from  $r$ -MaxRICS, but the same reduction introduced in [3] shows the following intractability when  $r = 3$ :

**Theorem 2 ([3]):** 3-MaxRICS is  $\mathcal{NP}$ -hard.

However, it would be hard to show the hardness of approximating  $r$ -MaxRICS for  $r \geq 3$  by using a similar reduction with small modification to the reduction in [3]. In this paper, by using a different *gap-preserving* reduction, we first show the following inapproximability of 3-MaxRICS.

**Theorem 3:** 3-MaxRICS cannot be approximated in polynomial time within a factor of  $n^{1/6-\varepsilon}$  for any constant  $\varepsilon > 0$  if  $\mathcal{P} \neq \mathcal{NP}$ , where  $n$  is the number of vertices in the input graph.

Furthermore, by using additional ideas to the reduction, we show the same inapproximability of  $r$ -MaxRICS for any fixed integer  $r \geq 4$ .

**Corollary 1:** For any fixed integer  $r \geq 4$ ,  $r$ -MaxRICS cannot be approximated in polynomial time within a factor of  $n^{1/6-\varepsilon}$  for any constant  $\varepsilon > 0$  if  $\mathcal{P} \neq \mathcal{NP}$ , where  $n$  is the number of vertices in the input graph.

The proofs of Theorem 3 and Corollary 1 will be given in Section 3.

## 1.2 Related Work

Recently, the problem of finding a maximum induced subgraph having regularity is very popular. Many researchers study the following variant, that is, the connectivity property is not imposed on the induced subgraph.

MAXIMUM  $r$ -REGULAR INDUCED SUBGRAPH ( $r$ -MaxRIS)

Input: A graph  $G = (V, E)$  and an integer  $r$ .

Goal: Find a maximum subset of vertices  $S \subseteq V$  such that the induced subgraph  $G[S]$  on  $S$  is  $r$ -regular.

If we do not require the connectivity constraint, then the problems when  $r = 0$  and  $r = 1$  correspond to the well studied MAXIMUM INDEPENDENT SET and MAXIMUM INDUCED MATCHING problems, respectively. The former problem is hard even to approximate [5]. The  $\mathcal{NP}$ -hardness of the latter problem is also shown in [1], [10]. In [9] Orlovich, Finke, Gordon, and Zverovich prove the MAXIMUM INDUCED MATCHING cannot be approximated within a factor of  $|V|^{1/2-\varepsilon}$  in polynomial time for any  $\varepsilon > 0$ . The parameterized complexity and exact exponential algorithms of  $r$ -MaxRIS are studied in [8] and [4], respectively. Very recently, in [2] Cardoso, Kamiński, and Lozin prove that  $r$ -MaxRIS is  $\mathcal{NP}$ -hard for any value of  $r \geq 3$ . Motivated by this result, we investigate the complexity of the connected version problem  $r$ -MaxRICS for  $r \geq 3$  in this paper.

## 2. Notation

By  $(u, v)$  we denote an edge with endpoints  $u$  and  $v$ . For a vertex  $u$ , the set of vertices adjacent to  $u$  in  $G$  is denoted by  $N_G(u)$  or simply by  $N(u)$ , and  $(u, N_G(u))$  denotes the set  $\{(u, v) \mid v \in N_G(u)\}$  of edges. Let the degree of a vertex  $u$  be denoted by  $\deg(u)$ , i.e.,  $|N(u)| = \deg(u)$ . A (simple) path  $P$  of length  $\ell$  from a vertex  $v_0$  to a vertex  $v_\ell$  is represented as a sequence of vertices such that  $P = \langle v_0, v_1, \dots, v_\ell \rangle$ , and  $|P|$  denotes the length of  $P$ . A cycle  $C$  of length  $\ell$  is similarly denoted by  $C = \langle v_0, v_1, \dots, v_{\ell-1}, v_0 \rangle$ , and  $|C|$  denotes the length of  $C$ . A *chord* of a path (cycle)  $\langle v_0, \dots, v_\ell \rangle$  ( $\langle v_0, \dots, v_{\ell-1}, v_0 \rangle$ ) is an edge between two vertices of the path (cycle) that is not an edge of the path (cycle). A path (cycle) is *chordless* if it contains no chords, i.e., an induced cycle must be chordless. Let  $G_1, G_2, \dots, G_\ell$  be  $\ell$  graphs and also let  $v_i$  be some vertex in  $G_i$  for  $1 \leq i \leq \ell$ . Then,  $\langle G_1, G_2, \dots, G_\ell \rangle$  denotes the subgraph  $G = (V(G_1) \cup V(G_2) \cup \dots \cup V(G_\ell), E(G_1) \cup E(G_2) \cup \dots \cup E(G_\ell) \cup \{(v_1, v_2), (v_2, v_3), \dots, (v_{\ell-1}, v_\ell)\})$ . That is, two adjacent graphs  $G_{i-1}$  and  $G_i$  are connected by only one edge and  $G$  roughly forms a path. Similarly,  $\langle G_1, G_2, \dots, G_\ell, G_1 \rangle$  roughly forms a cycle.

## 3. Hardness of Approximating $r$ -MaxRICS

In this section we give the proofs of Theorem 3 and Corollary 1. The hardness of approximating  $r$ -MaxRICS for  $r \geq 3$  is shown via a gap-preserving reduction from LONGEST INDUCED CYCLE problem, i.e., 2-MaxRICS. Consider an input graph  $G = (V(G), E(G))$  of 2-MaxRICS with  $n$  vertices and  $m$  edges. Then, we construct a graph  $H = (V(H), E(H))$  of  $r$ -MaxRICS. First we show the  $O(n^{1/6-\varepsilon})$ -inapproximability of 3-MaxRICS and then the same  $O(n^{1/6-\varepsilon})$ -inapproximability of the general  $r$ -MaxRICS problem for  $r \geq 4$ .

Let  $OPT_1(G)$  (and  $OPT_2(H)$ , respectively) denote the number of vertices of an optimal solution for  $G$  of 2-MaxRICS (and  $H$  of  $r$ -MaxRICS, respectively). Let  $V(G) = \{v_1, v_2, \dots, v_n\}$  of  $n$  vertices and  $E(G) = \{e_1, e_2, \dots, e_m\}$  of  $m$  edges. Let  $g(n)$  be a parameter function of the instance  $G$ . Then we provide the gap preserving reduction such that (C1) if  $OPT_1(G) \geq g(n)$ , then  $OPT_2(H) \geq 4(n^3 + 1) \times g(n)$ , and (C2) if  $OPT_1(G) < \frac{g(n)}{n^{1-\varepsilon}}$  for a positive constant  $\varepsilon$ , then  $OPT_2(H) < 4(n^3 + 1) \times \frac{g(n)}{n^{1-\varepsilon}}$ . As we will explain it, the number of vertices in the reduced graph  $H$  is  $O(n^6)$ . Hence the approximation gap is  $n^{1-\varepsilon} = O(|V(H)|^{1/6-\varepsilon})$  for any constant  $\varepsilon > 0$ .

### 3.1 Reduction for $r = 3$

Without loss of generality, we can assume that there is no vertex whose degree is one in the input graph  $G$




Figure 1: Subgraph  $H_i$

of 2-MaxRICS. The reason is that such a vertex does not contribute to any feasible solution, i.e., a cycle, of 2-MaxRICS and can be removed from  $G$ .

The constructed graph  $H$  consists of (i)  $n$  subgraphs,  $H_1$  through  $H_n$ , which are associated with  $n$  vertices,  $v_1$  through  $v_n$ , respectively, and (ii)  $m$  edge sets,  $E_1$  through  $E_m$ , which are associated with  $m$  edges,  $e_1$  through  $e_m$ , respectively.

(i) Here we describe the construction of the  $i$ th subgraph  $H_i$  in detail for some  $i$  ( $1 \leq i \leq n$ ). See Figure 1, which illustrates  $H_i$ . Suppose that the set of vertices adjacent to  $v_i$  is  $N(v_i) = \{v_{i_1}, v_{i_2}, \dots, v_{i_{deg(v_i)}}\}$ , where  $i_j \in \{1, 2, \dots, n\} \setminus \{i\}$  for  $1 \leq j \leq deg(v_i)$ . The subgraph  $H_i = (V(H_i), E(H_i))$  includes  $deg(v_i)$  vertices,  $u_{i, i_1}$  through  $u_{i, i_{deg(v_i)}}$  that correspond to the vertices adjacent to  $v_i$ , and  $deg(v_i)(deg(v_i) - 1)/2$  path gadgets,  $P_{i_1, i, i_2}, P_{i_1, i, i_3}, \dots, P_{i_1, i, i_{deg(v_i)}}, P_{i_2, i, i_3}, \dots, P_{i_{deg(v_i)-1}, i, i_{deg(v_i)}}$ , where two vertices  $u_{i, i_j}$  and  $u_{i, i_k}$  are connected via the path gadget  $P_{i_j, i, i_k}$  for  $v_{i_j}, v_{i_k} \in N(v_i)$ . As an example, in Figure 1, the top vertex  $u_{i, i_1}$  and the bottom  $u_{i, i_4}$  are connected via  $P_{i_1, i, i_4}$ . Each path gadget  $P_{i_j, i, i_k}$  includes  $n^3$  subgraphs,

$P_{i_j, i, i_k}^1$  through  $P_{i_j, i, i_k}^{n^3}$ , where, for each  $1 \leq p \leq n^3$ ,

$$V(P_{i_j, i, i_k}^p) = \{w_{i_j, i, i_k}^{p,1}, w_{i_j, i, i_k}^{p,2}, w_{i_j, i, i_k}^{p,3}, \gamma_{i_j, i, i_k}^p\},$$

$$E(P_{i_j, i, i_k}^p) = (\gamma_{i_j, i, i_k}^p, \{w_{i_j, i, i_k}^{p,1}, w_{i_j, i, i_k}^{p,2}, w_{i_j, i, i_k}^{p,3}\}) \cup \{(w_{i_j, i, i_k}^{p,1}, w_{i_j, i, i_k}^{p,2}), (w_{i_j, i, i_k}^{p,2}, w_{i_j, i, i_k}^{p,3})\}.$$

In the path gadget  $P_{i_j, i, i_k}$ , two vertices  $w_{i_j, i, i_k}^{p,1}$  and  $w_{i_j, i, i_k}^{p,3}$  are respectively identical to the vertices  $u_{i, i_j}$  and  $u_{i, i_k}$  prepared in the above. For  $2 \leq p \leq n^3$ , contiguous two subgraphs  $P_{i_j, i, i_k}^{p-1}$  and  $P_{i_j, i, i_k}^p$  are connected by one edge  $(w_{i_j, i, i_k}^{p-1,3}, w_{i_j, i, i_k}^{p,1})$  except for a pair  $P_{i_j, i, i_k}^{q-1}$  and  $P_{i_j, i, i_k}^q$  for some  $q$ : the two subgraphs  $P_{i_j, i, i_k}^{q-1}$  and  $P_{i_j, i, i_k}^q$  are connected by a path of length four  $\langle w_{i_j, i, i_k}^{q-1,3}, \beta_{i_j, i, i_k}^1, \beta_{i_j, i, i_k}^2, \beta_{i_j, i, i_k}^3, w_{i_j, i, i_k}^{q,1} \rangle$ . This  $q$  can be arbitrary since we just want to insert the path of length four into the path gadget, and as an example,  $q = 3$  in the path gadget  $P_{i_1, i, i_4}$  in Fig. 1. Finally, we prepare a special vertex  $\alpha_i$ , and  $\alpha_i$  is connected to all  $\{\beta_{i_1, i, i_4}^1, \beta_{i_1, i, i_4}^2, \beta_{i_1, i, i_4}^3\}$ 's. In the following,  $\alpha_1, \alpha_2, \dots, \alpha_n$  are called  $\alpha$ -vertices. Similarly,  $\beta$ -vertices and  $\gamma$ -vertices mean the vertices labeled




Figure 2:  $E_k$  connecting  $H_i$  and  $H_j$

by  $\beta$  and  $\gamma$ , respectively. Since each path gadget has  $4n^3 + 3$  vertices (two of which are shared with other path gadgets), the total number of vertices in  $H_i$  is

$$|V(H_i)| = \frac{\deg(v_i)(\deg(v_i) - 1)(4n^3 + 1)}{2} + n + 1,$$

i.e., there are  $O(n^5)$  vertices in  $H_i$ .

(ii) Next we explain construction of the edge sets  $E_1$  through  $E_m$ . Now suppose that  $e_k$  connects  $v_i$  with  $v_j$  for  $i \neq j$ . Also suppose that the sets of vertices adjacent to  $v_i$  and  $v_j$  are  $N(v_i) = \{j, i_2, \dots, i_{\deg(v_i)}\}$  and  $N(v_j) = \{i, j_2, \dots, j_{\deg(v_j)}\}$ , respectively. Then,  $(u_{i,j}, u_{j,i}) \in E_k$  where  $u_{i,j} \in V(H_i)$  in the  $i$ th subgraph  $H_i$  and  $u_{j,i} \in V(H_j)$  in the  $j$ th subgraph  $H_j$ . Furthermore, by the following rules,  $\gamma$ -vertices in the path gadgets are connected: See Figure 2. Every vertex in the path gadget  $P_{x,i,y}$  for  $x = j$  or  $y = j$  in  $H_i$  is not connected to any vertex in  $H_j$ , except for  $u_{i,j}$ . Similarly, every vertex in  $P_{s,j,t}$  for  $s = i$  or  $t = i$  in  $H_j$  is not connected to  $H_i$ , except for  $u_{j,i}$ . For a path gadget  $P_{x,i,y}$  in  $H_i$ , where  $j \notin \{x, y\}$  we prepare a set of edges as follows. Let  $D = \min_{k \in \{i, j\}} \{\deg(v_k)(\deg(v_k) - 1)/2 - (\deg(v_k) - 1)\}$ .

- In  $P_{x,i,y}$ , there are  $n^3$   $\gamma$ -vertices,  $\gamma_{x,i,y}^1$  through  $\gamma_{x,i,y}^{n^3}$ . Consider  $D$   $\gamma$ -vertices among those  $n^3$   $\gamma$ -vertices, the  $((j-1)n^2 + 1)$ th vertex  $\gamma_{x,i,y}^{(j-1)n^2+1}$  through the  $((j-1)n^2 + D)$ th vertex  $\gamma_{x,i,y}^{(j-1)n^2+D}$ .
- Next take a look at the  $j$ th subgraph  $H_j$  and the path gadgets  $P_{s,j,t}$ 's for  $i \notin \{s, t\}$ . Note that the number of such gadgets is  $\deg(v_j)(\deg(v_j) - 1)/2 - (\deg(v_j) - 1)$  and hence at least  $D$ . Then, consider the  $((i-1)n^2 + 1)$ th vertex  $\gamma_{s,j,t}^{(i-1)n^2+1}$  in each  $P_{s,j,t}$ . Here, the term

“+1” in the superscript of  $\gamma$  comes from the assumption that  $j_1 = i$ ; if  $j_k = i$ , we consider the  $((i-1)n^2 + k)$ th  $\gamma$ -vertex.

- Then, we can choose any function  $f$  which assigns each element in  $\{1, \dots, D\}$  to a string  $s, j, t$  such that  $i \notin \{s, t\}$  and it holds  $f(b) \neq f(c)$  if  $b \neq c$ . Finally, we connect  $\gamma_{x,i,y}^{(j-1)n^2+k}$  with  $\gamma_{f(k)}^{(i-1)n^2+1}$  for  $1 \leq k \leq D$ . It is important that the path gadget  $P_{x,i,y}$  is connected to  $P_{s,j,t}$  via only one edge.

Just to make the above construction clear, see Figure 3. For example, if an input instance  $G$  is the left graph, then the reduced graph  $H$  is illustrated in the right graph, where some details on the path gadgets are omitted due to the space. For example, since two vertices  $v_1$  and  $v_2$  are connected via the edge  $e_1$  in  $G$ ,  $u_{1,2}$  in  $H_1$  is connected to  $u_{2,1}$  in  $H_2$ . Similarly to  $e_2$  through  $e_6$ , there are five edges,  $(u_{1,3}, u_{3,1})$ ,  $(u_{3,4}, u_{4,3})$ ,  $(u_{2,4}, u_{4,2})$ ,  $(u_{2,5}, u_{5,2})$ , and  $(u_{4,5}, u_{5,4})$  in  $H$ . Furthermore, two path gadgets  $P_{1,2,5}$  and  $P_{3,4,5}$  are connected by one edge  $(\gamma_1, \gamma_2)$ .

Each subgraph  $H_i$  has  $O(n^5)$  vertices and thus the total number of vertices  $|V(H)| = O(n^6)$ . Clearly, this reduction can be done in polynomial time. In the next two subsections, we show that both conditions (C1) and (C2) are satisfied by the above reduction.

### 3.2 Proof of Condition (C1)

Without loss of generality, suppose that a longest induced cycle in  $G$  is  $C^* = \langle v_1, v_2, \dots, v_\ell, v_1 \rangle$  of length  $\ell$ , and thus  $OPT_1(G) = |C^*| = \ell \geq g(n)$ . Then we select the following subset  $S$  of  $4(n^3 + 1) \times \ell$  vertices and the induced




Figure 3: Input graph  $G$  (left) and reduced graph  $H$  (right)

subgraph  $G[S]$ :

$$S = V(P_{\ell,1,2}) \cup \{\alpha_1\} \cup V(P_{1,2,3}) \cup \{\alpha_2\} \\ \cup \dots \cup V(P_{\ell-1,\ell,1}) \cup \{\alpha_\ell\}.$$

For example, take a look at the graph  $G$  illustrated in Figure 3 again. One can see that the longest induced cycle in  $G$  is  $\langle v_1, v_3, v_4, v_2, v_1 \rangle$ . Then, we select the connected subgraph induced on the following set of vertices:

$$V(P_{2,1,3}) \cup \{\alpha_1\} \cup V(P_{1,3,4}) \cup \{\alpha_3\} \\ \cup V(P_{2,4,3}) \cup \{\alpha_4\} \cup V(P_{1,2,4}) \cup \{\alpha_2\}$$

It is easy to see that the induced subgraph is 3-regular and connected. Hence, the reduction satisfies the condition (C1).

### 3.3 Proof of Condition (C2)

We show that the reduction satisfies the condition (C2) by showing its contraposition. Suppose that  $OPT_2(H) \geq 4(n^3 + 1) \cdot \frac{g(n)}{n^{1-\varepsilon}}$  holds for a positive constant  $\varepsilon$ , and  $S^*$  is an optimal set of vertices such that the subgraph  $H[S^*]$  induced on  $S^*$  is connected and 3-regular. In the following, one of the crucial observations is that we can select at most one path gadget from each subgraph  $H_i$  into the optimal set  $S^*$  of vertices, and if a portion of the path gadget is only selected, then the induced subgraph cannot be 3-regular.

(I) See Figure 1 again. Suppose for example that two path gadgets  $P_{i_1,i,i_4}$  and  $P_{i_2,i,i_3}$  are selected, and put their vertices into  $S^*$ . In order to make the degree of  $\beta$ -vertices three, we need to also select  $\alpha_i$ . However, the degree of  $\alpha_1$  becomes six. This implies that we can select at most three  $\beta$ -vertices from each subgraph  $H_i$ .

(II) From the above observation (I), we consider the case that at most two of  $\beta_{j,i,k}^1$ ,  $\beta_{j,i,k}^2$ , and  $\beta_{j,i,k}^3$  are selected for some  $i, j, k$ . Let us assume that we select  $\beta_{j,i,k}^1$  and  $\beta_{j,i,k}^2$  ( $\beta_{j,i,k}^1$  and  $\beta_{j,i,k}^3$ , resp.) are put into  $S^*$ , but  $\beta_{j,i,k}^3$  ( $\beta_{j,i,k}^2$ , resp.) is not selected. Then, the degree of  $\beta_{j,i,k}^2$  ( $\beta_{j,i,k}^1$  and  $\beta_{j,i,k}^3$ , resp.) is at most 2 even if we select  $\alpha_i$ , i.e., the induced subgraph cannot be 3-regular. By a similar reason, we can not select only one of the  $\beta$ -vertices. Hence, if we select  $\beta$ -vertices, all of the three  $\beta$ -vertices in one path gadget must be selected.

As for  $w$ -vertices, a similar discussion can be done: For example, if we select  $w_{j,i,k}^{p,1}$  and  $w_{j,i,k}^{p,3}$  for some  $i, j, k, p$ , but  $w_{j,i,k}^{p,2}$  ( $\gamma_{j,i,k}^p$ , resp.) is not selected, then the degree of  $\gamma_{j,i,k}^p$  ( $w_{j,i,k}^{p,2}$ , resp.) is only 2. Thus, we need to select all the vertices of the part  $P_{k,i,j}^p$  if we select some vertices from it.

Combining two observations above, one can see that the edges connecting  $P_{k,i,j}^{p-1}$  and  $P_{k,i,j}^p$ , or  $w$ -vertices and  $\beta$ -vertices are necessary to make the degrees of the vertices three. As a result, we can conclude that if only a part of one path gadget is chosen, then the induced subgraph obtained




Figure 4: Modified path gadget in the proof of Corollary 1

cannot be 3-regular.

(III) From (I) and (II), we can assume that if some vertices of a path gadget are selected into  $S^*$ , it means that the whole vertices of the path gadget is selected. For example, suppose that  $P_{i_1, i_4}$  is selected. Since the degree of the endpoint  $u_{i_1}$  ( $u_{i_4}$ ) of  $P_{i_1, i_4}$  is only 2, we have to put at least one vertex into  $S^*$  from another subgraph adjacent to  $H_{i_1}$ , say, a vertex  $u_{j, i}$  in  $H_j$ . This implies that the induced subgraph  $H[S^*]$  forms a cycle-like structure  $\langle H_{i_1}, H_{i_2}, \dots, H_{i_j}, H_{i_1} \rangle$  connecting  $H_{i_1}, H_{i_2}, \dots, H_{i_j}, H_{i_1}$  in order, where  $\{i_1, i_2, \dots, i_j\} \subseteq \{1, 2, \dots, n\}$ .

We mention that such an induced subgraph  $H[S^*]$  is 3-regular if and only if the corresponding subgraph in the original graph  $G$  is an induced cycle. The if-part is clear by the discussion of the previous section. Let us look at the induced subgraph  $H[V(P_{2,1,3}) \cup V(P_{1,3,4}) \cup V(P_{3,4,5}) \cup V(P_{2,5,4}) \cup V(P_{1,2,5})]$  in the right graph  $H$  shown in Figure 3. Then, the induced subgraph includes the chord edge  $(\gamma_1, \gamma_2)$  and thus the degree of  $\gamma_1$  and  $\gamma_4$  is 4. The reason why the induced subgraph cannot be 3-regular comes from the fact that the cycle  $\langle v_1, v_3, v_4, v_5, v_2, v_1 \rangle$  includes the chord edge  $(v_1, v_4)$  in the original graph  $G$ . The edges between  $\gamma$ -vertices are placed because there is an edge between their corresponding vertices in  $G$ . As a result, the assumption that  $H[S^*]$  is an optimal solution, i.e., 3-regular, implies that the corresponding induced subgraph in the original graph  $G$  forms a cycle  $\langle v_{i_1}, v_{i_2}, \dots, v_{i_j}, v_{i_1} \rangle$ .

Since the number of vertices in each path gadget is  $4(n^3 + 1)$ ,  $OPT_1(G) \geq \frac{g(n)}{n^{1-\varepsilon}}$  holds by the assumption  $OPT_2(H) \geq 4(n^3 + 1) \cdot \frac{g(n)}{n^{1-\varepsilon}}$ . Therefore, the condition (C2) is also satisfied.

### 3.4 Reduction for $r \geq 4$

In this section, we give a brief sketch of the ideas to prove Corollary 1, i.e., the  $O(n^{1/6-\varepsilon})$  inapproximability for  $r$ -MaxRICS for any fixed integer  $r \geq 4$ .

The proof is very similar to that of Theorem 3. The main difference between those proofs is the structure of each path gadget. See Figure 4, which shows the modified

path gadget. (i) We replace each of  $\gamma$ -vertices in Figure 1 with the complete graph  $K_{r-2}$  of  $r - 2$  vertices, and then connect one  $\gamma$ -vertex in  $H_i$  and one  $\gamma$ -vertex in  $H_j$  for  $i \neq j$  by a similar manner to the reduction for the case  $r = 3$ . (ii) As for  $\beta$ -vertices, we prepare  $K_{r-2}$  of  $r - 2$  vertices, say,  $\beta^1, \dots, \beta^{r-2}$ , and two vertices, say,  $\beta^0$  and  $\beta^{r-1}$ , such that each of the two vertices  $\beta^0$  and  $\beta^{r-2}$  is adjacent to all the vertices in  $K_{r-2}$ . Then, all of the  $\beta$ -vertices are connected to the  $\alpha$ -vertex similar to the reduction for  $r = 3$ . Since the reduction requires  $n^3$   $\gamma$ -vertices to connect all the pairs of  $H_i$ 's, which is independent of the value of  $r$ , the path gadget consists of  $\lceil \frac{n^3}{r-2} \rceil$  subgraphs, say,  $P_{j,i,k}^1$  through  $P_{j,i,k}^{\lceil \frac{n^3}{r-2} \rceil}$ . Further details are omitted here.

### Acknowledgment

This work is partially supported by Grant-in-Aid for Scientific Research (KAKENHI), 22700019 and 23500020.

### References

- [1] K. Cameron. Induced Matchings. *Discrete Applied Math*, 24, pp.97–102 (1989)
- [2] D.M. Cardoso, M. Kamiński, and V. Lozin. Maximum  $k$ -regular induced subgraphs. *J. Combinatorial Optimization*, 14(5), pp.455–463 (2007)
- [3] V. Bonifaci, U. Di Iorio, and L. Laura. The complexity of uniform Nash equilibria and related regular subgraph problems. *Theoretical Computer Science*, 401, pp.144–152 (2008)
- [4] S. Gupta, V. Raman, and S. Saurabh. Fast exponential algorithms for maximum  $r$ -regular induced subgraph problems. In *Proc. FSTTCS 2006*, pp.139–151 (2006)
- [5] J. Håstad. Clique is hard to approximate within  $n^{1-\varepsilon}$ . *Acta Mathematica*, 182 (1), pp.105–142, 1999.
- [6] V. Kann. Strong lower bounds on the approximability of some NPO PB-complete maximization problems. In *Proc. MFCS 1995*, pp.227–236 (1995)
- [7] C. Lund and M. Yannakakis. The approximation of maximum subgraph problems. In *Proc. ICALP 1993*, pp.40–51 (1993)
- [8] H. Moser and S. Sikdar. The parameterized complexity of the induced matching problem. *Discrete Applied Mathematics*, 157, pp.715–727 (2009)
- [9] Y. Orlovich, G. Finke, V. Gordon, and I. Zverovich. Approximability results for the maximum and minimum maximal induced matching problems. *Discrete Optimization*, 5, pp.584–593 (2008)
- [10] L.J. Stockmeyer and V.V. Vazirani.  $\mathcal{NP}$ -completeness of some generalizations of the maximum matching problem. *Information Processing Letters*, 15(1), pp.14–19 (1982)